



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**A cretaceous genus of Abietineae.**—Dr. STOPES<sup>38</sup> has investigated a petri-fied tree trunk from the Cretaceous of New Zealand, which shows “a striking mixture of abietinean and araucarian characters.” She regards it as a new genus, which is named *Planoxylon*, and which is made to contain two species, the one investigated (*P. Hectori*) and another (*P. Lindleii*) that had been referred to *Araucarioxylon* by SEWARD. *P. Hectori* is of special interest because “it comes from Australasia, where no Abietineae or Juniperineae or other forms with abietinean ray structure are endemic at present.”—J. M. C.

**Morphology of Treubia.**—GRÜN,<sup>39</sup> studying material of *Treubia insignis* Goebel from Tjibodas, finds that the thallus develops from a dolabrate apical cell like most of the anacrogynous Jungermanniales; that the archegonia occur in clusters of 8–10 and may have as many as 16 neck canal cells each; that only one sporophyte develops on each thallus; that vegetative reproduction is by means of 3 or 4-celled brood bodies; that mycorrhiza are present in the thallus; and that the chromosome numbers are 8 and 16.—W. J. G. LAND.

**Plagiochasma.**—In a monograph of the genus *Plagiochasma*, EVANS<sup>40</sup> includes as much of the morphology of the group as will be useful for purposes of identification of species, gives a key to the North American species, and describes in detail the 6 species found in North America, one of which is new to science. The reviewer believes that several new species will be added to the group from Mexico when botanical exploration again becomes possible in that country.—W. J. G. LAND.

**Endemism in the flora of New York.**—TAYLOR<sup>41</sup> has found in the flora of the region near New York 22 endemic species, of which he considers 14 to be accounted for by generic or specific instability; 2 are “habitat endemics,” or habitat forms of more widely spread species; while 5 are evidently relic endemics. He is convinced, therefore, that many forces are operative in making endemics, and he gives some discussion of such possible forces.—GEO. D. FULLER.

---

<sup>38</sup> STOPES, MARIE C., An early type of the Abietineae(?) from the Cretaceous of New Zealand. *Ann. Botany* 30:111–125. *pl. 4. figs. 7.* 1916.

<sup>39</sup> GRÜN, C., Monographische Studien an *Treubia insignis* Goebel. *Flora* 106: 331–392. *pls. 3–5. figs. 14.* 1914.

<sup>40</sup> EVANS, ALEXANDER W., The genus *Plagiochasma* and its North American species. *Bull. Torr. Bot. Club* 42:259–308. *figs. 8.* 1915.

<sup>41</sup> TAYLOR, NORMAN, Endemism in the flora of the vicinity of New York. *Torreya* 16:18–27. 1916.